

Get the conversation going with your healthcare provider so you can reduce your tardive dyskinesia (TD) symptoms.

Answer the questions below, and bring this with you to your next appointment so you can share with your healthcare team the impact TD has on your life.

Once-daily AUSTEDO XR contains the same active ingredient as twice-daily AUSTEDO® (deutetrabenazine) tablets. AUSTEDO has not been demonstrated to affect all of the outcomes mentioned below.



## 1 Select the word(s) you might use to describe your symptoms.

Select all that apply.

- |  |   |  |  |
|--|---|--|--|
| <input type="checkbox"/> Foot/leg/hand movements | <input type="checkbox"/> Movements while sleeping | <input type="checkbox"/> Writhing or twitching           | <input type="checkbox"/> Fingers/toes moving       |
| <input type="checkbox"/> Eye blinking            | <input type="checkbox"/> Twisting tongue          | <input type="checkbox"/> Clenching/biting/grinding teeth | <input type="checkbox"/> Face tics or lip smacking |

## 2 Select the word(s) you might use to describe how your symptoms make you feel, physically.

Select all that apply.

- |  |   |                                  |                                   |
|--|---|----------------------------------|-----------------------------------|
| <input type="checkbox"/> Unwanted sensations | <input type="checkbox"/> Moving involuntarily | <input type="checkbox"/> Jittery | <input type="checkbox"/> Restless |
| <input type="checkbox"/> In pain             | <input type="checkbox"/> Uncontrolled         |                                  |                                   |

## 3 Select the word(s) you might use to describe how your symptoms make you feel, emotionally.

Select all that apply.

- |   |   |  |  |
|---|---|--|--|
| <input type="checkbox"/> I notice that small tasks are becoming more difficult. | <input type="checkbox"/> I'm embarrassed that people have mentioned my movements. | <input type="checkbox"/> I'm self-conscious and feel like everyone is looking at me. | <input type="checkbox"/> I want to avoid crowded places and isolate to avoid interactions with family and friends. |
| <input type="checkbox"/> I feel I have no control over my body.                 | <input type="checkbox"/> None of the above  |  |  |

Living with TD can be frustrating and overwhelming. Sometimes, it can be difficult to describe it in your own words. This real patient quote may give you inspiration for what to say to your healthcare provider.



### APPROVED USE

AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

### IMPORTANT SAFETY INFORMATION

**AUSTEDO XR and AUSTEDO can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not** start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Please read additional **Important Safety Information** continued on the next page and click [here](#) or visit [www.AUSTEDO.com](http://www.AUSTEDO.com) to read or print the **Medication Guide** for AUSTEDO XR.

Get the conversation going with your healthcare provider so you can reduce your tardive dyskinesia (TD) symptoms.

Answer the questions below, and bring this with you to your next appointment so you can share with your healthcare team the impact TD has on your life.

Once-daily AUSTEDO XR contains the same active ingredient as twice-daily AUSTEDO® (deutetrabenazine) tablets. AUSTEDO has not been demonstrated to affect all of the outcomes mentioned below.



## 4 Are you open to treating your TD with an additional medication?

Please select one.

Yes

Somewhat open

Not sure

## 5 Select which of these statement(s) best describe how you feel about your next appointment.

Select all that apply.

I'm nervous about what will happen when I bring up my movements.

I'm concerned I won't be taken seriously.

I'm embarrassed to talk about my movements.

I do not want to take another medication.

I'm concerned about insurance coverage or affording AUSTEDO XR.

I'm worried about potential side effects.

None of the above

Preparing for the conversation with your healthcare provider can help facilitate your discussion. Bring information about your mental health conditions and treatments, as well as details about how your movements are affecting you.

### IMPORTANT SAFETY INFORMATION (CONTINUED)

#### Do not take AUSTEDO XR or AUSTEDO if you:

- have Huntington's disease and are depressed or have thoughts of suicide.
- have liver problems.
- are taking reserpine. **Do not** take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.
- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. **Do not** start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.
- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of

tetrabenazine.

- are taking valbenazine.

#### Other possible serious side effects include:

- **Irregular heartbeat (QT prolongation).** AUSTEDO XR and AUSTEDO increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.
- **Neuroleptic Malignant Syndrome.** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, or increased sweating.
- **Restlessness.** You may get a condition where you feel a strong urge to move. This is called akathisia.
- **Parkinsonism.** Symptoms include: slight shaking, body stiffness, trouble moving, trouble keeping your balance, or falls.

Please read additional **Important Safety Information** continued on the next page and click [here](#) or visit [www.AUSTEDO.com](http://www.AUSTEDO.com) to read or print the **Medication Guide** for AUSTEDO XR.

## **AUSTEDO XR is a once-daily treatment for adults with tardive dyskinesia (TD).**

Once-daily AUSTEDO XR contains the same active ingredient as twice-daily AUSTEDO® (deutetrabenazine) tablets.

**Use these questions to talk to your healthcare provider about treating TD and to learn more about AUSTEDO XR.**

### **1 How may AUSTEDO XR be able to help me?**

---

---

### **2 I don't want to change my mental health medications. Can I take AUSTEDO XR with my other medications?**

---

---

### **3 I'm concerned about side effects. What are the potential side effects of AUSTEDO XR?**

---

---

### **4 AUSTEDO XR has different dosing options. What does this mean for my treatment?**

---

---

### **5 How do I take AUSTEDO XR?**

---

---

### **6 What is the Abnormal Involuntary Movement Scale (AIMS), and how can it measure my progress?**

---

---

### **7 What support and resources are available if I start taking AUSTEDO XR?**

---

---

## **IMPORTANT SAFETY INFORMATION (CONTINUED)**

**Sleepiness (sedation) is a common side effect of AUSTEDO XR and AUSTEDO.** While taking AUSTEDO XR or AUSTEDO, do not drive a car or operate dangerous machinery until you know how AUSTEDO XR or AUSTEDO affects you. Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO XR or AUSTEDO may increase any sleepiness caused by AUSTEDO XR and AUSTEDO.

**The most common side effects of AUSTEDO in people with Huntington's disease include** sleepiness (sedation), diarrhea, tiredness, and dry mouth.

**The most common side effects of AUSTEDO in people with tardive dyskinesia include** inflammation of the nose and throat (nasopharyngitis) and problems sleeping (insomnia).

The most common side effects of AUSTEDO XR are expected to be similar to AUSTEDO in people with Huntington's disease or tardive dyskinesia.

These are not all the possible side effects of AUSTEDO XR or AUSTEDO. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call **1-800-FDA-1088**.

Please read the **Important Safety Information** and click [here](#) or visit [www.AUSTEDO.com](http://www.AUSTEDO.com) to read or print the **Medication Guide** for AUSTEDO XR.