

For the treatment of **tardive dyskinesia (TD)** in adults

ONCE-DAILY
Austedo® XR
(deutetrabenazine)
extended-release
6 mg, 12 mg, 18 mg, 24 mg, 30 mg,
36 mg, 42 mg, and 48 mg tablets

TD TREATMENT INFORMATION

*Reduce TD movement with
AUSTEDO XR—personalized
dosing that can be taken with
most mental health meds*



AS YOU GO WITH AUSTEDO



APPROVED USE

AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat the movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

IMPORTANT SAFETY INFORMATION

AUSTEDO XR and AUSTEDO® (deutetrabenazine) tablets can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Please read **Important Safety Information** on pages 8-9 and click [here](#) or visit www.AUSTEDO.com to read or print the **Medication Guide** for AUSTEDO XR.

teva

What is tardive dyskinesia (TD)?

SYMPTOMS AND CAUSES OF TD

TD is repetitive and uncontrollable movements in the face, arms, legs, and trunk that appear as twitching, shaking, or jerking. **TD symptoms may appear as:**



Flicking the fingers



Flailing arms or legs



Shrugging shoulders



Grimacing



Twisting or jerking



Irregular eye movements or blinking

Certain prescription medications used to treat mental health or gastrointestinal conditions can cause TD. Tardive dyskinesia is more than a side effect of using these medications—**it's a real condition with its own diagnosis and treatment plan.**



Medications that can cause TD include antipsychotics, antidepressants, and antiemetics.

If you're taking these medications and experiencing uncontrollable movements, ask your healthcare provider to evaluate you for TD

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not take AUSTEDO® XR (deutetrabenazine) extended-release tablets or AUSTEDO® (deutetrabenazine) tablets if you:

- have Huntington's disease and are depressed or have thoughts of suicide.
- have liver problems.

What is tardive dyskinesia (TD)?

THE IMPACT OF TD

The unintentional and uncontrollable movements of TD:

- ✓ Can be mistaken for being drunk or on drugs
- ✓ Make it difficult to speak clearly
- ✓ Attract unwanted attention when you're out in public
- ✓ Affect your ability to read or write
- ✓ Make cutting your food or holding your drink challenging



TD is not your fault and can be treated.

Talk to your healthcare provider about
AUSTEDO® XR (deutetrabenazine) extended-release tablets.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not take AUSTEDO XR or AUSTEDO® (deutetrabenazine) tablets if you:

- are taking reserpine. **Do not** take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.

About treatment

BENEFITS OF AUSTEDO® XR (DEUTETRABENAZINE) EXTENDED-RELEASE TABLETS



AUSTEDO XR reduces TD movement with **one once-daily, extended-release pill.***



With AUSTEDO XR, you don't have to choose between maintaining your mental health and treating TD. **AUSTEDO XR can be taken with most mental health medications.**



Personalized dosing with AUSTEDO XR allows you and your healthcare provider to work together to find a dose that's effective for you.



Scan this QR code or visit www.AUSTEDO.com to view the common side effects of AUSTEDO XR.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not take AUSTEDO XR or AUSTEDO® (deutetrabenazine) tablets if you:

- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. **Do not** start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.

About treatment

In a 12-week clinical study,
some people experienced
>2X REDUCTION
in TD movements vs placebo.[†]

Some people saw
a response as early as
2 WEEKS.

In a separate study,
established safety results
~3 YEARS.[‡]



Cost shouldn't stand in the way of treatment.
Teva **Shared Solutions**[®] will verify coverage and work through
financial assistance options with you to help make treatment affordable

*Once-daily AUSTEDO[®] XR (deutetrabenazine) extended-release tablets contains the same active ingredient as twice-daily AUSTEDO[®] (deutetrabenazine) tablets. Data on this page is based on twice-daily dosing.

[†]AUSTEDO reduced the Abnormal Involuntary Movement Scale (AIMS) total score by 3.3 points vs 1.4 points with placebo. For people taking AUSTEDO, this represents a 33% reduction in AIMS total score vs a 14% reduction for those taking placebo.

[‡]This was an open-label study, which is a drug trial in which all participants are aware of the drug being tested.

IMPORTANT SAFETY INFORMATION (CONTINUED)


Do not take AUSTEDO[®] XR (deutetrabenazine) extended-release tablets or AUSTEDO[®] (deutetrabenazine) tablets if you:

- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of tetrabenazine.
- are taking valbenazine.

How to get started

A PERSONALIZED APPROACH TO DOSING

Your healthcare provider will start you at a low dose of AUSTEDO® XR (deutetrabenazine) extended-release tablets and gradually increase your dose over time. This approach allows your body to get used to the medication and can help with side effects as movements are reduced. This way, your doctor can find a dose that works for you. This process is called titration.



Pills are for illustration only and not actual size or likeness.

Starting dose

Your doctor will start you on a low dose of AUSTEDO XR and will work with you to find your maintenance dose.

Maintenance dose

Once you reach a dose that provides effective movement reduction and is tolerated, that will become your maintenance dose. If you want to increase your dose, you and your doctor can discuss more dosing options.



Because everyone is different, your maintenance dose will be the one that works specifically for you.



IMPORTANT SAFETY INFORMATION (CONTINUED)

Other possible serious side effects include:

- **Irregular heartbeat (QT prolongation).** AUSTEDO XR and AUSTEDO® (deutetrabenazine) tablets increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.

How to get started

HOW TO TAKE YOUR MEDICATION



Once daily



With water



Whole (do not
crush or break)



With or
without food

If you cannot swallow AUSTEDO® XR (deutetrabenazine) extended-release tablets/AUSTEDO® (deutetrabenazine) tablets whole, tell your healthcare provider. You may need a different medication.

HOW YOU'LL BEGIN TREATMENT



Not actual image of the Sample Titration Kit.

Your doctor may provide 1 of these convenient options:

A Sample Titration
Kit to take home

— or —

A Prescription Titration
Kit to pick up

The Titration Kit will guide you through your first month with AUSTEDO XR.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Other possible serious side effects include:

- **Neuroleptic Malignant Syndrome.** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, or increased sweating.

APPROVED USE

AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat the movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

IMPORTANT SAFETY INFORMATION

AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Do not take AUSTEDO XR or AUSTEDO if you:

- have Huntington's disease and are depressed or have thoughts of suicide.
- have liver problems.
- are taking reserpine. **Do not** take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.
- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. **Do not** start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.
- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of tetrabenazine.
- are taking valbenazine.

IMPORTANT SAFETY INFORMATION (CONTINUED)**Other possible serious side effects include:**

- **Irregular heartbeat (QT prolongation).** AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.
- **Neuroleptic Malignant Syndrome.** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, or increased sweating.
- **Restlessness.** You may get a condition where you feel a strong urge to move. This is called akathisia.
- **Parkinsonism.** Symptoms include: slight shaking, body stiffness, trouble moving, trouble keeping your balance, or falls.

Sleepiness (sedation) is a common side effect of AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets.

While taking AUSTEDO XR or AUSTEDO, do not drive a car or operate dangerous machinery until you know how AUSTEDO XR or AUSTEDO affects you. Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO XR or AUSTEDO may increase any sleepiness caused by AUSTEDO XR and AUSTEDO.

The most common side effects of AUSTEDO in people with Huntington's disease include sleepiness (sedation), diarrhea, tiredness, and dry mouth.

The most common side effects of AUSTEDO in people with tardive dyskinesia include inflammation of the nose and throat (nasopharyngitis) and problems sleeping (insomnia).

The most common side effects of AUSTEDO XR are expected to be similar to AUSTEDO in people with Huntington's disease or tardive dyskinesia.

These are not all the possible side effects of AUSTEDO XR or AUSTEDO. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

Please read the accompanying **Medication Guide** [here](#).

Teva Shared Solutions® patient support program

REACH OUT TO US, WE'LL REACH FOR SOLUTIONS TOGETHER

With **Shared Solutions**, you'll have the benefit of a Patient Access Specialist for insurance and pharmacy questions, and nurse support for treatment and dosing guidance. Your **Shared Solutions** team are dedicated professionals in their field, so you'll get real help from real people.

Shared Solutions helps with financial, insurance, and prescription assistance so you can receive treatment sooner and stay on treatment

FINANCIAL SUPPORT

Whether you're insured, underinsured, or not insured at all, **Shared Solutions** will verify coverage and work through financial assistance options to help make treatment affordable. Plus, we'll help you navigate the paperwork so nothing falls through the cracks.



Close to 90% of people taking AUSTEDO® (deutetrabenazine) tablets pay \$10 or less per month for their prescription.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Other possible serious side effects include:


- **Irregular heartbeat (QT prolongation).** AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.

Teva Shared Solutions® patient support program

NURSE SUPPORT

While taking AUSTEDO® XR (deutetrabenazine) extended-release tablets with support from Teva **Shared Solutions**, you will receive calls throughout treatment from our nurses to see how we can best support you in reaching your goals. They can help you with:

- ✓ Learning about TD and AUSTEDO XR
- ✓ Starting a treatment plan, reaching your appropriate AUSTEDO XR dose, and even managing your refills
- ✓ Taking and remembering to take your AUSTEDO XR
- ✓ Providing tools and resources to help you stay on track



teva | Shared Solutions
Start your enrollment today!



1-800-887-8100

Monday-Friday
from 8 AM to 8 PM CT

Or visit [MySharedSolutions.com](https://www.MySharedSolutions.com)

IMPORTANT SAFETY INFORMATION (CONTINUED)

Other possible serious side effects include:

- **Neuroleptic Malignant Syndrome.** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, or increased sweating.



START TREATMENT. REDUCE TD MOVEMENTS.

- **More than 2x movement reduction at 12 weeks vs placebo^{††}**
- **One pill, once-daily extended release**
- **Can be safely taken with other mental health medications**
- **Start at no cost**
- **Get support through Teva Shared Solutions[®] patient support program**



Scan this code

to access other resources. Escanee este código para acceder a otros recursos.

*Once-daily AUSTEDO[®] XR (deutetrabenzina) extended-release tablets contains the same active ingredient as twice-daily AUSTEDO[®] (deutetrabenzina) tablets. Data on this page is based on twice-daily dosing.

^{††}AUSTEDO reduced the AIMS total score by 3.3 points vs 1.4 points with placebo. For participants taking AUSTEDO, this represents a 33% reduction in AIMS total score vs a 14% reduction for those taking placebo.

IMPORTANT SAFETY INFORMATION

AUSTEDO XR and AUSTEDO[®] (deutetrabenzina) tablets can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Please read **Important Safety Information** on pages 8-9 and click [here](#) or visit www.AUSTEDO.com to read or print the **Medication Guide** for AUSTEDO XR.