


  
**UZEDY**<sup>TM</sup>
  
 (risperidone)

50 mg 75 mg 100 mg 125 mg

150 mg 200 mg 250 mg

extended-release  
injectable suspension



For adults living with schizophrenia,

**RELAPSE PREVENTION WITH UZEDY.\***

**INNER STRENGTH BY YOU.**

\*During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.

## IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including **Boxed WARNING**.

# SCHIZOPHRENIA IS COMPLEX

Schizophrenia affects everyone differently.

It can be unpredictable and impact many areas of life. Trying to treat your schizophrenia can be a challenge too.

Medication is an important part of treating schizophrenia. But getting offtrack with your medication schedule could lead to a setback or even a relapse.

**That's why finding a treatment that's right for you is so important.**

# UZEDY MAY BE ABLE TO HELP

When it comes to treating your schizophrenia, do you feel you're making progress? If you're looking for a different schizophrenia medication, it may be time to think about UZEDY.



With an injection every 1 or 2 months, UZEDY is a long-acting treatment option for schizophrenia that doesn't require a daily pill.



UZEDY uses a syringe with a short needle that injects a small amount of medicine.\*

\*UZEDY is supplied as a prefilled syringe designed to provide 0.14 to 0.7 mL of medication, depending on the dose, with a 5/8-inch needle.

## IMPORTANT SAFETY INFORMATION (CONTINUED)


**Do not receive UZEDY (risperidone) extended-release injectable suspension if you** are allergic to risperidone, paliperidone, or any of its components.

**UZEDY may cause serious side effects, including:**

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including **Boxed WARNING**.**



  
**UZEDY**<sup>TM</sup>  
(risperidone) extended-release  
injectable suspension

50 mg 75 mg 100 mg 125 mg  
150 mg 200 mg 250 mg

# MORE STABILITY, FEWER SYMPTOMS\*

UZEDY can help you reduce your risk of relapse and maintain stability with your symptoms.\*

## IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including:

- **Uncontrolled facial or body movements (tardive dyskinesia)** that may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including **Boxed WARNING**.

\*In a clinical study of UZEDY, stability was defined based on patients who stayed outpatient, had few symptoms from a list of specific psychotic symptoms, and had symptom and clinical functioning scores below certain cutoffs for at least 4 weeks.



# WHAT MAKES UZEDY DIFFERENT? IT'S IN THE DETAILS

UZEDY was researched in a clinical study that included 542 patients who were treated with either UZEDY or placebo. **This study was the largest and longest pivotal trial for a risperidone long-acting injectable in schizophrenia to date.**



## IN A CLINICAL STUDY†

**UZEDY reduced the risk of relapse by up to 80% compared with placebo.**

Of the patients receiving once-monthly UZEDY, 7% relapsed vs 29% of patients receiving placebo.



## IN A SURVEY‡

**92% of patients with schizophrenia (58 out of 63) were satisfied with UZEDY.**

Patients surveyed received UZEDY as part of a clinical trial.

## DID YOU KNOW?

**Risperidone is one of the most frequently prescribed medications proven to treat schizophrenia.**

†During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.

‡Data were collected from 63 patients, 24 physicians, and 25 nurses in a prospective, cross-sectional companion survey assessing the perceptions regarding ease of use and satisfaction with UZEDY. The survey was administered after a minimum of 2 experiences prescribing, administering, or receiving UZEDY.

## IMPORTANT SAFETY INFORMATION (CONTINUED)

### UZEDY may cause serious side effects, including:

- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including **Boxed WARNING**.**



### FEWER SETBACKS

With once-monthly UZEDY, **the relapse-free rate of patients was 93% after 24 weeks (about 6 months) of treatment**; 72% was the relapse-free rate of patients receiving placebo.



### GREATER STABILITY

In the clinical study, **87% of patients maintained stability** with once-monthly UZEDY; 61% of patients receiving placebo did so.



### IMPROVED SYMPTOMS

In the clinical study, patients who received UZEDY showed **continued improvement in their schizophrenia symptoms**.



### SAFETY WAS EXTENSIVELY STUDIED

**The most common side effects of risperidone included** slow movements, stiffness, shaking, restlessness, abnormal muscle contractions or movements, drowsiness, dizziness, anxiety, blurred vision, nausea, vomiting, indigestion, diarrhea, increased saliva, constipation, dry mouth, increased appetite, weight gain, tiredness, rash, and common cold symptoms. Injection site reactions including a lump or itching were reported with UZEDY.

## IMPORTANT SAFETY INFORMATION (CONTINUED)

**UZEDY may cause serious side effects, including:**

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including **Boxed WARNING**.**



# WHAT TO EXPECT



## UZEDY USES A SYRINGE WITH A SHORT NEEDLE THAT INJECTS A SMALL AMOUNT OF MEDICINE\*

In fact, it's one of the smallest LAI needles used to treat schizophrenia. In a survey, **7 out of 10 patients felt UZEDY was a better injection experience than other LAIs they'd taken in the past.**<sup>†</sup>



## UZEDY ALSO HELPS YOU STREAMLINE YOUR START TO TREATMENT

That means no extra risperidone pills or additional starter doses once you start UZEDY. Plus, you could go up to 2 months between injections. When given the choice to keep taking UZEDY or return to their previous medication, **90% of surveyed patients (57 out of 63) chose to continue their treatment with UZEDY.**<sup>†</sup>

### IMPORTANT SAFETY INFORMATION (CONTINUED)

#### UZEDY may cause serious side effects, including:

- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including **Boxed WARNING**.**



\*UZEDY is supplied as a prefilled syringe designed to provide 0.14 to 0.7 mL of medication, depending on dose, with a 5/8-inch needle.

<sup>†</sup>Data were collected from 63 patients, 24 physicians, and 25 nurses in a prospective, cross-sectional companion survey assessing the perceptions regarding ease of use and satisfaction with UZEDY. The survey was administered after a minimum of 2 experiences prescribing, administering, or receiving UZEDY.

# A TREATMENT WITH OPTIONS

## 1- OR 2-MONTH DOSING

Instead of taking a pill every day to treat your schizophrenia symptoms, with UZEDY you get 1 injection from your healthcare provider every 1 or 2 months.



1 INJECTION  
EVERY MONTH

OR



1 INJECTION  
EVERY 2 MONTHS

## STAYING ON SCHEDULE

Once you and your healthcare provider decide UZEDY is right for you, it's important to stay on your treatment plan. Schedule appointments every 1 to 2 months, depending on your dosing schedule.

## MULTIPLE ADMINISTRATION SITES

UPPER ARM OR ABDOMEN


## IMPORTANT SAFETY INFORMATION (CONTINUED)

**UZEDY may cause serious side effects, including:**

- **Low white blood cell count.**
- **Problems thinking clearly and moving your body.** Do not drive, operate machinery, or do other dangerous activities until you know how UZEDY affects you.
- **Seizures (convulsions).**
- **Difficulty swallowing that can cause food or liquid to get into your lungs.**
- **Prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including **Boxed WARNING**.**





**UZEDY**<sup>TM</sup>  
(risperidone)

50 mg 75 mg 100 mg 125 mg

150 mg 200 mg 250 mg

extended-release  
injectable suspension



**89%**

**IN A SURVEY\***

**89% of patients (56 out of 63) said receiving UZEDY was easy.**

\*Data were collected from 63 patients, 24 physicians, and 25 nurses in a prospective, cross-sectional companion survey assessing the perceptions regarding ease of use and satisfaction with UZEDY. The survey was administered after a minimum of 2 experiences prescribing, administering, or receiving UZEDY.

## **IMPORTANT SAFETY INFORMATION (CONTINUED)**

**UZEDY may cause serious side effects, including:**

- **Problems with control of your body temperature (too high or too low). Avoid getting overheated or dehydrated.**

**The most common side effects of risperidone included** slow movements, stiffness, shaking, restlessness, abnormal muscle contractions or movements, drowsiness, dizziness, anxiety, blurred vision, nausea, vomiting, indigestion, diarrhea, increased saliva, constipation, dry mouth, increased appetite, weight gain, tiredness, rash, and common cold symptoms. Injection site reactions including a lump or itching were reported with UZEDY. These are not all the possible side effects of UZEDY. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.

**Do not** drink alcohol during treatment with UZEDY.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.**


# FINANCIAL ASSISTANCE AND ONGOING SUPPORT

**teva** | Shared Solutions

Teva Shared Solutions<sup>®</sup> is a support program to help you plan for starting and staying on UZEDY. We help streamline your start to treatment by helping you—or a caregiver—figure out some of the details. Shared Solutions can:

- ✓ Confirm prescription coverage and pharmacy options based on eligibility and coverage
- ✓ Explain Medicare and Medicaid coverage options
- ✓ See if you qualify for the Patient Assistance Program (PAP), which provides free product to eligible patients
- ✓ Help you understand prior authorization requirements
- ✓ Work with you to find an injection site near your home or work
- ✓ Provide over-the-phone nurse support to answer questions and help you stay on UZEDY





## YOU COULD SAVE ON UZEDY



With the Savings Offer for UZEDY, eligible patients may pay as little as \$0 per dose.\*

To get started, call 1-800-887-8100 (Monday-Friday, 9 AM-8 PM ET) or go to [UZEDY.com](http://UZEDY.com).

\*Offer is available for patients with commercial insurance only. This offer is NOT available for patients eligible for Medicare, Medicaid, or any other form of government insurance coverage. Please see complete Terms and Conditions at [www.UZEDY.com](http://www.UZEDY.com).

The Patient Savings Program for UZEDY™ (risperidone) extended-release injectable suspension is available to eligible patients who have been prescribed UZEDY and have commercial prescription insurance. Eligible patients may pay as little as \$0 for once-monthly or once-every-2-month dosing options of UZEDY. Maximum program assistance per prescription and annual benefit limits per individual apply and out-of-pocket expenses may vary. Patient is responsible for costs above maximum benefit amounts. This program is restricted to residents of the United States and United States territories. Uninsured and cash-paying patients are NOT eligible for this program. Patients enrolled in any state or federally funded healthcare program, including but not limited to, Medicare, Medigap, Medicaid, VA, DOD, TRICARE, Puerto Rico Government Health Insurance Plan, Medicare-eligible patients enrolled in an employer-sponsored health plan or prescription drug benefit program for retirees, are NOT eligible for this Program. Teva Neuroscience, Inc. and its affiliates reserve the right to change, rescind, revoke, or discontinue this program at any time without notice. Please see complete Terms and Conditions at [www.UZEDY.com](http://www.UZEDY.com).

### IMPORTANT SAFETY INFORMATION (CONTINUED)

**Before receiving UZEDY, tell your healthcare provider** about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome.
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia).
- have diabetes or have a family history of diabetes.
- have had dizziness or fainting or are being treated for high blood pressure.
- have had a low white blood cell count.
- have or have had seizures or epilepsy.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.**

# TALK TO YOUR HEALTHCARE PROVIDER ABOUT UZEDY

Talking about your schizophrenia and treatment options can be tough. That's why we've included information to help you start a conversation with your healthcare provider.

## BE SURE TO ASK THESE QUESTIONS AT YOUR NEXT APPOINTMENT:

- ? Is UZEDY an option for me?
- ? What are possible benefits and risks with UZEDY?
- ? What are the most common side effects with UZEDY?
- ? How can UZEDY fit into my treatment routine?

## IMPORTANT SAFETY INFORMATION (CONTINUED)

**Before receiving UZEDY (risperidone) extended-release injectable suspension, tell your healthcare provider** about all your medical conditions, including if you:

- are pregnant or plan to become pregnant during treatment with UZEDY. It is not known if UZEDY will harm your unborn baby. Use of UZEDY during the third trimester of pregnancy may cause side effects in the newborn infant, including agitation, abnormal muscle tone, tremor, drowsiness, difficulty feeding, and difficulty breathing. Seek medical attention if you notice these signs. If you become pregnant during treatment with UZEDY, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics, or call 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.**





**UZEDY**<sup>TM</sup>  
(risperidone) extended-release  
injectable suspension

**SHARE THIS INFORMATION  
WITH YOUR HEALTHCARE  
PROVIDER, AS WELL:**

- ✓ All your current symptoms
- ✓ Any medications you're taking now
- ✓ Any diagnosis you've received in the past
- ✓ Your family's health history

**92%**

**IN A SURVEY\***

**92% of patients with schizophrenia  
(58 out of 63) were satisfied with  
UZEDY.**

## **IMPORTANT SAFETY INFORMATION (CONTINUED)**

**Before receiving UZEDY (risperidone) extended-release injectable suspension, tell your healthcare provider** about all your medical conditions, including if you:

- are breastfeeding or plan to breastfeed. If you are receiving UZEDY and are breastfeeding, monitor your infant for sleepiness, inadequate weight gain, jitteriness, tremors, and abnormal muscle movements. Seek medical care if you notice these signs.
- have or have had kidney or liver problems.

**Tell your healthcare provider about all the medicines you take or plan to take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. UZEDY and other medicines may affect each other.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.**

\*Data were collected from 63 patients, 24 physicians, and 25 nurses in a prospective, cross-sectional companion survey assessing the perceptions regarding ease of use and satisfaction with UZEDY. The survey was administered after a minimum of 2 experiences prescribing, administering, or receiving UZEDY.



## APPROVED USE

UZEDY (risperidone) extended-release injectable suspension is a prescription medicine used to treat schizophrenia in adults.

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about UZEDY?

**UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.**


**Do not receive UZEDY if you** are allergic to risperidone, paliperidone, or any of its components.

### UZEDY may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.
- **Uncontrolled facial or body movements (tardive dyskinesia)** that may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.
- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes

(e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.
- **Low white blood cell count.**
- **Problems thinking clearly and moving your body.** Do not drive, operate machinery, or do other dangerous activities until you know how UZEDY affects you.
- **Seizures (convulsions).**
- **Difficulty swallowing that can cause food or liquid to get into your lungs.**
- **Prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.
- **Problems with control of your body temperature (too high or too low). Avoid getting overheated or dehydrated.**



**The most common side effects of risperidone included** slow movements, stiffness, shaking, restlessness, abnormal muscle contractions or movements, drowsiness, dizziness, anxiety, blurred vision, nausea, vomiting, indigestion, diarrhea, increased saliva, constipation, dry mouth, increased appetite, weight gain, tiredness, rash, and common cold symptoms. Injection site reactions including a lump or itching were reported with UZEDY. These are not all the possible side effects of UZEDY. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.

**Do not** drink alcohol during treatment with UZEDY.

**Before receiving UZEDY, tell your healthcare provider** about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome.
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia).
- have diabetes or have a family history of diabetes.
- have had dizziness or fainting or are being treated for high blood pressure.
- have had a low white blood cell count.
- have or have had seizures or epilepsy.
- are pregnant or plan to become pregnant during treatment with UZEDY. It is not known if UZEDY will harm your unborn baby. Use of UZEDY during the third trimester of pregnancy may cause side effects in the newborn infant, including agitation, abnormal muscle tone, tremor, drowsiness, difficulty feeding, and difficulty breathing. Seek medical attention if

you notice these signs. If you become pregnant during treatment with UZEDY, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics, or call 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.

- are breastfeeding or plan to breastfeed. If you are receiving UZEDY and are breastfeeding, monitor your infant for sleepiness, inadequate weight gain, jitteriness, tremors, and abnormal muscle movements. Seek medical care if you notice these signs.
- have or have had kidney or liver problems.

**Tell your healthcare provider about all the medicines you take or plan to take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. UZEDY and other medicines may affect each other.

**You are encouraged to report side effects of prescription drugs to the FDA. Visit** [www.fda.gov/medwatch](http://www.fda.gov/medwatch), **or call** 1-800-FDA-1088.

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.**

**UZEDY**<sup>TM</sup>  
(risperidone) extended-release  
injectable suspension

50 mg 75 mg 100 mg 125 mg  
150 mg 200 mg 250 mg



## READY TO GET STARTED?

Talk to your doctor if you think UZEDY might be right for you.  
Visit [UZEDY.com](https://www.uzedy.com) to learn more and find support and financial assistance.

### APPROVED USE

UZEDY (risperidone) extended-release injectable suspension is a prescription medicine used to treat schizophrenia in adults.

### IMPORTANT SAFETY INFORMATION

#### What is the most important information I should know about UZEDY?

**UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.**

**Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.**



© 2023 Teva Neuroscience, Inc.  
RIS-40649 September 2023