


teva | Shared Solutions

# SERVICES + OFFERINGS

FOR ONGOING SUPPORT



**UZEDY**<sup>TM</sup>  
(risperidone) extended-release  
injectable suspension

50 mg 75 mg 100 mg 125 mg  
150 mg 200 mg 250 mg

## INDICATION AND USAGE

UZEDY (risperidone) extended-release injectable suspension for subcutaneous use is indicated for the treatment of schizophrenia in adults.

## IMPORTANT SAFETY INFORMATION

### WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. UZEDY is not approved for use in patients with dementia-related psychosis and has not been studied in this patient population.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING.



# LET'S WORK TOGETHER



Teva Shared Solutions® is designed and committed to help patients gain affordable access to UZEDY.

A dedicated team provides support through the following services and offerings to help patients get started and stay on treatment.



## Patient Initiation and Coordination

Help patients get started with UZEDY

### **BENEFITS VERIFICATION**

Confirms prescription coverage and pharmacy options based on specific eligibility and coverage

### **PRIOR AUTHORIZATION/APEALS SUPPORT**

Communicates the prior authorization requirements, and supports the appeals process as requested

### **MEDICARE AND MEDICAID BENEFITS NAVIGATION SUPPORT**

Reviews Medicare and Medicaid coverage options

### **COORDINATION WITH A DISPENSING PHARMACY**

Coordinates care with the patient, prescriber, dispensing pharmacy, and site of care



## Financial Assistance

Help patients identify financial support options for UZEDY

### PATIENT ASSISTANCE PROGRAM

Provides free product to eligible patients

### SAVINGS OFFER

Reduces costs for commercially insured patients (eligible patients may pay as little as <del>\$0</del> for once-monthly or once-every-2-month dosing options of UZEDY)\*

\*Offer is available for patients with commercial insurance only. This offer is NOT available for patients eligible for Medicare, Medicaid, or any other form of government insurance coverage.



## Alternate Site-of-Care Network

Help patients find convenient site-of-care locations

### DIRECTORY

Provides a directory of available treatment locations



## Nurse Support

Help patients stay informed about their treatment journey with UZEDY

### OVER-THE-PHONE SUPPORT AND EDUCATION

Provides an introductory program welcome call and nurse support to patients and caregivers to answer questions and help with treatment adherence

#### ADDITIONAL OFFERING

## Hospital Inpatient Free Trial Program for UZEDY

Provides UZEDY to eligible inpatient hospitals that cannot accept Prescription Drug Marketing Act-compliant samples. Visit [UZEDYHCP.com](http://UZEDYHCP.com) to review enrollment criteria.



Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including Boxed WARNING.



## INDICATION AND USAGE

UZEDY (risperidone) extended-release injectable suspension for subcutaneous use is indicated for the treatment of schizophrenia in adults.

## IMPORTANT SAFETY INFORMATION

### WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

**Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. UZEDY is not approved for use in patients with dementia-related psychosis and has not been studied in this patient population.**

**CONTRAINDICATIONS:** UZEDY is contraindicated in patients with a known hypersensitivity to risperidone, its metabolite, paliperidone, or to any of its components. Hypersensitivity reactions, including anaphylactic reactions and angioedema, have been reported in patients treated with risperidone or paliperidone.

### WARNINGS AND PRECAUTIONS

**Cerebrovascular Adverse Reactions:** In trials of elderly patients with dementia-related psychosis, there was a significantly higher incidence of cerebrovascular adverse events (e.g., stroke, transient ischemic attack), including fatalities, in patients treated with oral risperidone compared to placebo. UZEDY is not approved for use in patients with dementia-related psychosis.

**Neuroleptic Malignant Syndrome (NMS):** NMS, a potentially fatal symptom complex, has been reported in association with antipsychotic drugs. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status including delirium, and autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure. If NMS is suspected, immediately discontinue UZEDY and provide symptomatic treatment and monitoring.

**Tardive Dyskinesia (TD):** TD, a syndrome consisting of potentially irreversible, involuntary, dyskinetic movements, may develop in patients treated with antipsychotic drugs. Although the prevalence of the syndrome appears to be highest among the elderly, especially elderly women, it is impossible to predict which patients will develop the syndrome. Whether antipsychotic drug products differ in their potential to cause TD is unknown.

The risk of developing TD and the likelihood that it will become irreversible are believed to increase with the duration of treatment and the cumulative dose. The syndrome can develop, after relatively brief treatment periods, even at low doses. It may also occur after discontinuation. TD may remit, partially or completely, if antipsychotic treatment is discontinued. Antipsychotic treatment, itself, however, may suppress (or partially suppress) the signs and symptoms of the syndrome, possibly masking the underlying process. The effect that symptomatic suppression has upon the long-term course of the syndrome is unknown.

If signs and symptoms of TD appear in a patient treated with UZEDY, drug discontinuation should be considered. However, some patients may require treatment with UZEDY despite the presence of the syndrome. In patients who do require chronic treatment, use the lowest dose and the shortest duration of treatment producing a satisfactory clinical response. Periodically reassess the need for continued treatment.

**Metabolic Changes:** Atypical antipsychotic drugs have been associated with metabolic changes that may increase cardiovascular/cerebrovascular risk. These metabolic changes include hyperglycemia, dyslipidemia, and body weight gain. While all of the drugs in the class have been shown to produce some metabolic changes, each drug has its own specific risk profile.

**Hyperglycemia and diabetes mellitus (DM),** in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, have been reported in patients treated with atypical antipsychotics, including risperidone. Patients with an established diagnosis of DM who are started on atypical antipsychotics, including UZEDY, should be monitored regularly for worsening of glucose control. Patients with risk factors for DM (e.g., obesity, family history of diabetes) who are starting treatment with atypical antipsychotics, including UZEDY, should undergo fasting blood glucose (FBG) testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics, including UZEDY, should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics, including UZEDY, should undergo FBG testing. In some cases, hyperglycemia has resolved when the atypical antipsychotic, including risperidone, was discontinued; however, some patients required continuation of antidiabetic treatment despite discontinuation of risperidone.

**Dyslipidemia** has been observed in patients treated with atypical antipsychotics.

**Weight gain** has been observed with atypical antipsychotic use. Monitoring weight is recommended.

**Hyperprolactinemia:** As with other drugs that antagonize dopamine  $D_2$  receptors, risperidone elevates prolactin levels and the elevation persists during chronic administration. Risperidone is associated with higher levels of prolactin elevation than other antipsychotic agents.

**Orthostatic Hypotension and Syncope:** UZEDY may induce orthostatic hypotension associated with dizziness, tachycardia, and in some patients, syncope. UZEDY should be used with particular caution in patients with known cardiovascular disease, cerebrovascular disease, and conditions which would predispose patients to hypotension and in the elderly and patients with renal or hepatic impairment. Monitoring of orthostatic vital signs should be considered in all such patients, and a dose reduction should be considered if hypotension occurs. Clinically significant hypotension has been observed with concomitant use of oral risperidone and antihypertensive medication.

**Falls:** Antipsychotics, including UZEDY, may cause somnolence, postural hypotension, motor and sensory instability, which may lead to falls and, consequently, fractures or other fall-related injuries. Somnolence, postural hypotension, motor and sensory instability have been reported with the use of risperidone. For patients, particularly the elderly, with diseases, conditions, or medications that could exacerbate these effects, assess the risk of falls when initiating antipsychotic treatment and recurrently for patients on long-term antipsychotic therapy.

**Leukopenia, Neutropenia, and Agranulocytosis** have been reported with antipsychotic agents, including risperidone. In patients with a pre-existing history of a clinically significant low white blood cell count (WBC) or absolute neutrophil count (ANC) or a history of drug-induced leukopenia or neutropenia, perform a complete blood count (CBC) frequently during the first few months of therapy. In such patients, consider discontinuation of UZEDY at the first sign of a clinically significant decline in WBC in the absence of other causative factors. Monitor patients with clinically significant neutropenia for fever or other symptoms or signs of infection and treat promptly if such symptoms or signs occur. Discontinue UZEDY in patients with ANC < 1000/mm<sup>3</sup> and follow their WBC until recovery.

**Potential for Cognitive and Motor Impairment:** UZEDY, like other antipsychotics, may cause somnolence and has the potential to impair judgement, thinking, and motor skills. Somnolence was a commonly reported adverse reaction associated with oral risperidone treatment. Caution patients about operating hazardous machinery, including motor vehicles, until they are reasonably certain that treatment with UZEDY does not affect them adversely.

**Seizures** During premarketing studies of oral risperidone in adult patients with schizophrenia, seizures occurred in 0.3% of patients (9 out of 2,607 patients), two in association with hyponatremia. Use UZEDY cautiously in patients with a history of seizures or other conditions that potentially lower the seizure threshold.

**Dysphagia:** Esophageal dysmotility and aspiration have been associated with antipsychotic drug use. Antipsychotic drugs, including UZEDY, should be used cautiously in patients at risk for aspiration.

**Priapism** has been reported during postmarketing surveillance for other risperidone products. A case of priapism was reported in premarket studies of UZEDY. Severe priapism may require surgical intervention.

(continued on next page)



**Body temperature regulation.** Disruption of the body's ability to reduce core body temperature has been attributed to antipsychotic agents. Both hyperthermia and hypothermia have been reported in association with oral risperidone use. Strenuous exercise, exposure to extreme heat, dehydration, and anticholinergic medications may contribute to an elevation in core body temperature; use UZEDY with caution in patients who experience these conditions.

#### ADVERSE REACTIONS

The most common adverse reactions with risperidone ( $\geq 5\%$  and greater than placebo) were parkinsonism, akathisia, dystonia, tremor, sedation, dizziness, anxiety, blurred vision, nausea, vomiting, upper abdominal pain, stomach discomfort, dyspepsia, diarrhea, salivary hypersecretion, constipation, dry mouth, increased appetite, increased weight, fatigue, rash, nasal congestion, upper respiratory tract infection, nasopharyngitis, and pharyngolaryngeal pain.

The most common injection site reactions with UZEDY ( $\geq 5\%$  and greater than placebo) were pruritus and nodule.

#### DRUG INTERACTIONS

- Carbamazepine and other strong CYP3A4 inducers decrease plasma concentrations of risperidone.
- Fluoxetine, paroxetine, and other strong CYP2D6 inhibitors increase risperidone plasma concentration.
- Due to additive pharmacologic effects, the concomitant use of centrally-acting drugs, including alcohol, may increase nervous system disorders.
- UZEDY may enhance the hypotensive effects of other therapeutic agents with this potential.
- UZEDY may antagonize the pharmacologic effects of dopamine agonists.
- Concomitant use with methylphenidate, when there is change in dosage of either medication, may increase the risk of extrapyramidal symptoms (EPS)

#### USE IN SPECIFIC POPULATIONS

**Pregnancy:** May cause EPS and/or withdrawal symptoms in neonates with third trimester exposure. There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to atypical antipsychotics, including UZEDY, during pregnancy. Healthcare providers are encouraged to register patients by contacting the National Pregnancy Registry for Atypical Antipsychotics at 1-866-961-2388 or online at <http://womensmentalhealth.org/clinicaland-research-programs/pregnancyregistry/>.

**Lactation:** Infants exposed to risperidone through breastmilk should be monitored for excess sedation, failure to thrive, jitteriness, and EPS.


**Fertility:** UZEDY may cause a reversible reduction in fertility in females.

**Pediatric Use:** Safety and effectiveness of UZEDY have not been established in pediatric patients.

**Renal or Hepatic Impairment:** Carefully titrate on oral risperidone up to at least 2 mg daily before initiating treatment with UZEDY.

**Patients with Parkinson's disease or dementia with Lewy bodies** can experience increased sensitivity to UZEDY. Manifestations and features are consistent with NMS.

**Please click here for full [Prescribing Information](#), including Boxed WARNING, for UZEDY.**



**UZEDY**<sup>TM</sup>  
(risperidone) extended-release  
injectable suspension

50 mg 75 mg 100 mg 125 mg  
150 mg 200 mg 250 mg

SUPPORT FOR PATIENTS RECEIVING UZEDY  
AND THEIR CAREGIVERS

**teva** | Shared Solutions

**TO ENROLL**

**CALL 1-800-887-8100 (M-F 9 AM TO 8 PM ET)**

**VISIT [UZEDYHCP.com](https://www.uzedyhcp.com)**

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including **Boxed WARNING**.